

Fourier Mukai And Nahm Transforms In Geometry And Mathematical Physics

Fourier Mukai And Nahm Transforms In Geometry And Mathematical Physics

Summary:

Fourier Mukai And Nahm Transforms In Geometry And Mathematical Physics Download Textbook Pdf hosted by Skye Sawyer on October 21 2018. It is a file download of Fourier Mukai And Nahm Transforms In Geometry And Mathematical Physics that you can be grabbed it with no registration on texivoire.org. For your info, this site dont upload pdf download Fourier Mukai And Nahm Transforms In Geometry And Mathematical Physics on texivoire.org, it's just PDF generator result for the preview.

Fourier-Mukai transform - Wikipedia In algebraic geometry, a Fourier-Mukai transform \hat{K} is a functor between derived categories of coherent sheaves $D(X) \rightarrow D(Y)$ for schemes X and Y , which is, in a sense, an integral transform along a kernel object $K \in D(X \times Y)$. FOURIER-MUKAI PARTNERS OF SURFACES IN POSITIVE CHARACTERISTIC FOURIER-MUKAI PARTNERS OF K3 SURFACES IN POSITIVE CHARACTERISTIC MAX LIEBLICH AND MARTIN OLSSON CONTENTS 1. Introduction 2. Mukai motive 3. Kernels of Fourier-Mukai equivalences 9. big picture - Heuristic behind the Fourier-Mukai transform ... The Fourier-Mukai transform in algebraic geometry gets its name because it at least superficially resembles the classical Fourier transform. (And of course because it was studied by Mukai.) Let me give a rough picture of the Fourier-Mukai transform and how it resembles the classical situation.

Fourier-Mukai transforms for quotient varieties ... A Fourier-Mukai (FM) transform is an exact equivalence $\hat{K}: D(Y) \rightarrow D(X)$ between the bounded derived categories of coherent sheaves on two smooth projective varieties X and Y . FOURIER-MUKAI PARTNERS OF K3 SURFACES IN POSITIVE ... fourier-mukai partners of k3 surfaces in positive characteristic 3 of the appendix is Theorem A.1 concerning the Picard group of the general deformation of a fixed K3 surface from characteristic p to characteristic 0. Fourier-Mukai transform on abelian surfaces | SpringerLink We study moduli spaces of stable sheaves on abelian surfaces whose Mukai vectors are related by a cohomological Fourier-Mukai transform. We show that there is a Fourier-Mukai transform inducing a birational map between them.

Fourier-Mukai duality for K3 surfaces via Bridgeland ... Fourier-Mukai duality is a duality between a variety X and a moduli space of stable sheaves on X , which is a generalization of the duality between an abelian variety X and its dual abelian variety $\text{Pic}^0(X)$. In this article, we shall explain Fourier-Mukai duality for a K3 surface by using Bridgeland stability condition. GV-sheaves, Fourier-Mukai transform, and generic vanishing GV-SHEAVES, FOURIER-MUKAI TRANSFORM, AND GENERIC VANISHING By GIUSEPPE PARESCHI and MIHNEA POPA Abstract. We prove a formal criterion for generic vanishing, in the sense originated by Green.

fourier mukai transform